

# 3<sup>rd</sup> INTERNATIONAL OBESITY SUMMIT AND EXPO

&

2<sup>nd</sup> International Conference on

# DIABETES, NUTRITION, METABOLISM & MEDICARE

&

World Conference on

# LASER, OPTICS AND PHOTONICS

November 05 - 06, 2018 | Philadelphia, USA

Ida Fidelis E Denosta et al., Biomed Res 2018, Volume 29 | DOI: 10.4066/biomedicalresearch-C7-019

## MATERNAL RISK FACTORS ASSOCIATED WITH RETINOPATHY OF PREMATURITY AMONG PATIENTS IN EAST AVENUE MEDICAL CENTER FROM 2010 TO 2015

**Ida Fidelis E Denosta and Carina Cruz-Quimbo**

East Avenue Medical Center, Philippines

**Objectives:** This study aims to determine the maternal risk factors associated with the development of retinopathy of prematurity (ROP).

**Design:** A retrospective cohort study.

**Setting:** The study was done in a tertiary government hospital.

**Patients:** This study included admitted premature neonates in neonatal intensive care unit (NICU) who underwent ROP screening. All patients with congenital abnormality of the eyes were excluded in the study.

**Methodology:** List of admitted premature patients who underwent ROP screening was collected within the study period. Demographic data was collected from the patient's medical records.

**Statistical Analysis:** Descriptive statistics were presented for categorical outcome measures. Wilcoxon rank sum test was used to compare two averages. Chi-square test, Yates' chi-square test and Fisher's exact test were used to compare proportions. Univariate and multivariate regression analyses were used to estimate crude and adjusted odds-ratios, corresponding 95% CI and based on p-value  $\leq 0.05$  statistical significance.

**Result:** This study showed 57% cases of ROP, majority with Stage 1 (66%), mostly female infants (53.4%;  $p < 0.001$ ), mildly asphyxiated (51.2%,  $p = 0.290$ ; 77.2%,  $p = 0.117$ ) and were small for gestational age (57.3%;  $p < 0.001$ ). On the average, these infants underwent 1 day of either invasive or non-invasive mechanical ventilation, 2 days with FiO<sub>2</sub> of  $> 0.5$  liters per minute, had two blood transfusions, and with shorter hospital stay (17 vs 26 days;  $p < 0.001$ ). The average age of mothers who delivered infants with ROP was 26 years old, multipara (59.2%), and underwent normal spontaneous delivery (67%). About half reported adequate prenatal check-ups (53.8%), and diagnosed with urinary tract infection (40.1%).

**Conclusion:** Data showed that mothers with urinary tract infection and pre-eclampsia are more likely to deliver infants with retinopathy of prematurity thus significant maternal risk factors of ROP.

**Keywords:** Retinopathy of Prematurity, Neonates, Prematurity

## BIOGRAPHY

Ida Fidelis E Denosta has completed her degree of bachelor of science major in psychology at the age of 20 in Far Eastern University. She graduated as doctor of medicine at Fatima University Medical Center. She completed her residency training in pediatrics at east avenue medical center last December 2016 as an assistant chief resident. She is an active member of the Philippine Medical Association, Manila Medical Society and currently affiliated with Allied Care Experts (ACE) Medical Center-Quezon City and Caloocan City North Medical Center (CCNMC) as a junior consultant in the department of pediatrics.

[idadenosta@yahoo.com](mailto:idadenosta@yahoo.com)